

Surrey & Sussex gastroenteritis outbreak pack for education and childcare settings

The following guidance aims to help you manage an outbreak of gastroenteritis in an educational setting or nursery environment. The principles are effective at limiting the spread of viral and bacterial infections, including norovirus. It has been produced with reference to national guidance, entitled 'Health Protection in children and young people settings, including education' and can be found here - https://www.gov.uk/government/publications/health-protection-in-schools-and-otherchildcare-facilities

Contents

The guidance consists of the following sections:

- 1. Summary General principles for effective control of gastroenteritis
- Action Card 1 Key actions for the Head Teacher or Administrator to manage the outbreak.
- 3. Action Card 2 Key actions for caretakers and staff who will be undertaking cleaning, which is a vital step to reduce the spread of infection.
- 4. Action Card 3 Information to be provided to the Health Protection Team to help them give appropriate advice to manage the outbreak.
- 5. Action Card 4 Contact information for organisations that can provide help during an outbreak.
- 6. Links to the chapters of the national guidance Health Protection in educational settings and other childcare facilities. This includes the exclusion table and posters.
- 7. Information Sheet to send to parents/carers
- 8. Handwashing posters (general and for young children)

Summary for Norovirus Outbreaks				
Transmission Route:		Person to person by the faecal oral route, from aerosols of projectile vomit and from environmental contamination.		
Incubation Period:		12 to 48 hours (length of time from acquiring the infection to developing symptoms).		
Exclusion:		48 hours after their last episode of diarrhoea or vomiting.		
Closures:		It is not necessary to close the educational setting, unless there are operational reasons such as significant staff absence, which would be a decision for the educational setting in conjunction with the relevant Local Authority.		
Recommendations for action				
Hand Hygiene:	and after going to the toilet. Hand washing with liquid soap and warm water. Paper towels should be used for drying hands and a wastepaper bin provided for disposal. Alcohol gel is not effective against norovirus, but it can be used in addition to soap and water. Everyone should wash their hands-on arrival at the educational setting and before leaving.			
Cleaning and disinfection:	deternal det	are the educational setting surfaces are thoroughly cleaned daily using warm water and regent. ating surfaces and toilet areas should be disinfected after cleaning twice daily using a suchlorite solution 1,000 parts per million (such as Milton Solution) recommended that toilets are cleaned at least twice a day and their condition monitored age the day, with a view to an extra clean if required. responsible apron and gloves when cleaning. Dispose after use and wash hands as recommended that toilet including rims, seats, and handles All areas of the toilet including rims, seats, and handles Wash basins and taps Door handles and light switches All frequently touched equipment Frequently handled items such as telephones and computer keyboards Carpeted areas and soft furnishings should be cleaned as normal, preferably with a carpet shampoo rather than a vacuum cleaner (hypochlorite will bleach carpets and soft furnishings).		
Public spillage incident: Food Safety:	Cover Alway should After I disinfe is visil Carpe sham cleani The a	lages of vomit or diarrhoea occur: the excreta/vomit spillages immediately with disposable paper towels. It was wear a disposable apron and gloves when disposing of faeces/vomit. A face mask do be worn if there is a concern about splash contamination to the face. It removing the spillage, clean the surrounding area with warm soapy water, followed by ection with a hypochlorite solution of 1000 parts per million. Always clean a wider area than only contaminated. It is contaminated with faeces or vomit should be cleaned with warm soapy water (or a carpet poo) after removal of the spillage with paper towels. This should be followed by steaming if possible. It is should be followed by steaming if possible. It is should be followed by steaming if possible. It is should be followed by steaming if possible. It is should be followed by steaming if possible. It is should be followed by steaming if possible. It is should be followed by steaming if possible.		
Food Safety:	Oper easil Food orde	h boxes of chocolates/biscuits and fruit bowls must be removed in an outbreak as they can y become contaminated. I handlers should avoid contact with the educational setting children during an outbreak, in r to avoid contamination of the kitchen or food. Hens MUST have their own separate cleaning equipment.		

Further Information can be found in the guidance: Health Protection in children and young people settings, including education https://www.gov.uk/government/publications/health-protection-in-schools-and-other-childcare-facilities

Title Managing an outbreak of diarrhoea & vomiting in a school or

childcare setting

Description Overview of what to do in an outbreak situation

Who Those members of staff that are responsible for overseeing/managing an

outbreak i.e. Head Teacher, administrative staff, absence monitoring

manager etc.

Definitions of an outbreak

• An incident in which 2 or more people experiencing a similar illness are linked in time or place. (This is important for infections where there could be serious outcomes such as meningitis or measles)

• A greater than expected rate of infection compared with the usual background rate for the place and time where the outbreak has occurred (this is more applicable to gastro-intestinal or respiratory infections)

NOTE: You do not need to notify the Health Protection Team of all outbreaks of gastrointestinal infections routinely (see Action card 3).

What do you need to do?

- Ensure good record keeping names, dates of birth, symptoms, dates of onset of illness of cases, number and location of episodes of vomiting in communal areas etc. in the educational setting.
- Inform UK Health Security Agency (UKHSA) for advice & guidance (see Action Card 3 for triggers of when to notify).
- Prompt exclusion / isolation of affected children and staff, ensuring strict compliance with 48hour exclusion (i.e. fully recovered for 48 hours before returning to the educational setting)
- Consideration should be given to advising parents and visitors about the outbreak. An
 information sheet with advice is attached in section 7.
- Remind children about the importance of good hand hygiene.
- Cookery activities for the children as well as sand, playdough and water play activities should be suspended for the duration of the outbreak. All potentially contaminated materials should be discarded and replaced in accordance with your local procedure.
- Ensure robust infection control measures are in place:
 - Cleaning regime in place (see Action Card 2)
 - Guidance for the management of 'Spillages of vomit or diarrhoea' are followed (see Action Card 2)
 - Effective hand hygiene (see Summary Sheet)
 - Plentiful supply of personal protective equipment (PPE), liquid soap, paper towels, etc.
 - ➤ Non-essential visitors should be restricted and/or discouraged from visiting during the outbreak.
 - ➤ Dispose of all infection-exposed food items from communal areas i.e. boxes of chocolates, biscuits, or bowls of fruit.
- Declare the outbreak over when there has been 48 hours since the resolution of symptoms in the last known case.
- Terminal clean (Action card 2) to be done as soon as possible after resolution of outbreak.

Further Information can be found in the guidance: Health Protection in children and young people settings, including education https://www.gov.uk/government/publications/health-protection-in-schools-and-other-childcare-facilities

Title Cleaning & disinfection of the educational setting/nursery environment

Description Germs that cause diarrhoea & vomiting will survive in the environment, so to

stop the infection spreading it is important that the areas are maintained in a clean condition. Decontamination (cleaning and disinfecting) the areas

thoroughly on a frequent basis is necessary.

Who Caretakers, other staff with responsibility for cleaning

The following are terms that you may hear:

<u>Detergent:</u> A cleaning agent that helps to remove dirt and grease from surfaces

Disinfectant: A chemical that destroys bacteria.

Recommended Disinfectant: Chlorine - releasing agent/hypochlorite solution (e.g. bleach or 'Milton' solution 0.1% at 1000ppm)

Routine cleaning: A process that physically removes contamination which is done on a regular/daily basis. Warm water and detergent should be used to clean.

<u>Disinfection:</u> A process that reduces the number of germs to a level at which they are not harmful but is only effective if the surfaces and equipment are cleaned thoroughly with detergent and water beforehand.

Enhanced cleaning: During an outbreak cleaning and disinfection should be done twice daily as a minimum (plus as necessary). Particular attention should be given to frequently touched surfaces, e.g. door handles, flushes, taps, contact points, switches, mirrors, bins, furniture, stair rails, etc.

<u>Deep cleaning:</u> Intense and enhanced thorough cleaning process. Usually planned ahead e.g. during holiday periods.

<u>Terminal cleaning:</u> Cleaning that takes place at the end of an outbreak to ensure the area has been thoroughly cleaned and disinfected. This includes the steam cleaning of carpets and soft furnishings.

To clean effectively you will need:

Equipment

Clean, disposable, single-use cloths, dedicated mop & mop bucket (different coloured equipment for different areas is a useful tool), dustpan and brush, clinical waste bags (yellow), spillage kit, paper towels, Personal Protective Equipment (PPE), steam cleaner.

Cleaning equipment used by the kitchen staff **must** be kept separate and used solely for kitchen.

Chemicals

Detergent (see above)

Disinfectant (see above)

Sanitiser – this is a commercial purchased combination of a detergent with disinfectant properties.

You must always ensure that you follow your health and safety guidelines when handling chemicals including the use of appropriate PPE

- During an outbreak cleaning and disinfection should be done twice daily.
- Warm water and detergent should be used to clean hard surfaces followed by disinfection with a
 0.1% hypochlorite solution. Bleach or Milton are the recommended disinfectants (at 1000ppm),
 as they will kill both bacteria and viruses. If these are unsuitable, a disinfectant that has <u>both</u>
 antibacterial and antiviral properties **must** be used.
- Particular attention should be paid to cleaning and disinfecting toilet seats, toilet flush handles, door handles, washbasin taps, push-plates on doors, light switches, stair handrails and other frequently touched areas.
- Spillages of vomit or diarrhoea should be cordoned off and the area cleared immediately:
 - Wear disposable gloves and apron.
 - Use a spill kit or paper towels to soak up the spillage. Transfer these and any solid matter directly into a clinical waste bag.
 - Clean the soiled area with detergent and hot water, using a disposable cloth. A two-metre area around the spillage should be cleaned.
 - The same area should then be disinfected with the freshly made hypochlorite solution.
 - o Dispose of gloves, apron, and cloths into the clinical waste bag.
- Wash hands thoroughly using soap and water and dry them with paper towels. Alcohol hand-rub is not a substitute for hand washing after cleaning up a spillage.
- Vacuum cleaning carpets and floor buffing are not recommended during an outbreak.
- Do-not use bleach on carpets or soft furnishings a steam cleaner must be used to remove the bacterial or viral particles.
- Clothing contaminated with body fluids should be placed in a sealed plastic bag and taken home by the child/parent. Do not wash/sluice or soak the contaminated items on behalf of the child/parent.
- Carpets and soft furnishings should be steam-cleaned, or a carpet shampoo used.

Further Information can be found in the guidance: Health Protection in children and young people settings, including education https://www.gov.uk/government/publications/health-protection-in-educational settings-and-other-childcare-facilities/preventing-and-controlling-infections

Title Reporting an outbreak of D&V to UK Health Security Agency South East (UKHSA

SE) by telephone or email.

Description This card gives you the contact details for UKHSA SE and the information that the

UKHSA SE will require to help you manage your outbreak.

Who Those members of staff that are responsible for informing UKHSA SE of an

outbreak i.e. head teacher, administrative staff.

When to report:

Educational settings should notify the HPT in the following circumstances:

 Pupils and staff with vomiting and/or diarrhoea with onset of illness around the same time.

- Cases of vomiting and diarrhoea associated with an educational setting trip or event.
- Any severe gastrointestinal illness that has resulted in hospitalisation etc.
- Reports of severe or bloody diarrhoea.
- Increased levels of concern in parents/public/media.

The general increase in GI cases which occur periodically (which are usually caused by norovirus) can be managed by the educational setting using this guidance.

Phone: 0344 225 3861 or email se.acuteresponse@ukhsa.gov.uk

What UKHSA SE will ask for:

- Name and address of the educational setting, including the postcode
- Contact person's name and phone number
- Total number of children at the educational setting
- Details of the year groups / classes
- Number of staff at the educational setting
- Number of children affected, which year groups, details of siblings affected
- Number of staff affected
- Date of onset of first illness (onset date)
- Details of any recent educational setting trips, special events
- Details of any episodes of public vomiting
- Details of educational setting caterers affected
- Educational setting lunch / packed lunch ratio
- Educational setting layout and facilities, particularly who shares toilets

Further Information can be found in the guidance: Health Protection in children and young people settings, including education https://www.gov.uk/government/publications/health-protection-in-schools-and-other-childcare-facilities

Title Contacts and further information

Description This card provides details of the main contacts who can help you during

an outbreak.

WhoThose members of staff who are responsible for informing UKHSA SE

of an outbreak, i.e. administrative staff, head teacher

UKHSA: se.acuteresponse@ukhsa.gov.uk 0344 225 3861

Education Departments and Public Health Teams

Surrey County Council West Sussex County Council: public.health@surreycc.gov.uk In hours: 01243 642104

Out of Hours: 02031622286

East Sussex County Council Brighton and Hove City Council:

public.health@eastsussex.gov.uk <u>educationandskills@brighton-hove.gov.uk</u>

publichealth@brighton-hove.gov.uk

School Nursing Teams

	School nursing team: Telephone: 01273 696011 ext.1692 Email: sc- tr.adminbhschoolnurses@nhs.net	
	www.healthforkids.co.uk/sussex/meet-the-school-nurse-team-2/	
Brighton & Hove and West Sussex	Health Visiting teams: Telephone: 01273 266 000 Email: sc-tr.bh-healthvisitors@nhs.net	
	www.sussexcommunity.nhs.uk/services/healthy-child-programme-hcp-health-visitors-and-school-nurses-0-19-service/528135	
Surrey	www.childrenshealthsurrey.nhs.uk/services/school-nursing-general	
East Sussex	www.eastsussexchildren.nhs.uk/about/school-health	

Environmental Health Departments

Surrey EHOs:

Elmbridge foodteam@elmbridge.gov.uk

Epsom and Ewell ehpublichealth@epsom-ewell.gov.uk

Guildford regulatoryservices@guildford.gov.uk

Mole Valley env.health@molevalley.gov.uk

Reigate and Banstead infectious.disease@reigate-banstead.gov.uk

Runneymede environmentalhealth@runnymede.gov.uk

Spelthorne eh.commercial@spelthorne.gov.uk

Surrey Heath environmental.health@surreyheath.gov.uk

Tandridge eh@tandridge.gov.uk

Waverley environmentalhealth@waverley.gov.uk

Woking environmental.health@woking.gov.uk

Sussex EHOs:

Adur and Worthing id@adur-worthing.gov.uk

Arun infectious.disease@arun.gov.uk

Brighton and Hove ids@brighton-hove.gov.uk

Chichester healthprotection@chichester.gov.uk

Crawley environmentalservices@crawley.gov.uk

Hastings environmentalhealth@hastings.gov.uk

Horsham publichealth.licensing@horsham.gov.uk

Lewes and Eastbourne ehealth.ldc@lewes-eastbourne.gov.uk

Mid Sussex foodsafety@midsussex.gov.uk

Rother and Wealden envhealth@rother.gov.uk

UK Health Security Agency SE

Gastrointestinal Outbreak Pack for Education and Childcare Settings

National guidance is available on GOV UK website entitled 'Health protection in children and young people settings, including education'. The following are links to the individual chapters for ease of reading.

1.	What infections are, how they are transmitted and those at higher risk of infection	https://www.gov.uk/government/publications/health- protection-in-schools-and-other-childcare- facilities/what-infections-are-how-they-are- transmitted-and-those-at-higher-risk-of-infection
2.	Prevention and Controlling infections	https://www.gov.uk/government/publications/health- protection-in-schools-and-other-childcare- facilities/preventing-and-controlling-infections
3.	Supporting immunisation programmes	https://www.gov.uk/government/publications/health- protection-in-schools-and-other-childcare- facilities/supporting-immunisation-programmes
4.	Managing outbreaks and incidents	https://www.gov.uk/government/publications/health- protection-in-schools-and-other-childcare- facilities/managing-outbreaks-and-incidents
5.	Managing specific infectious diseases: A to Z	https://www.gov.uk/government/publications/health- protection-in-schools-and-other-childcare- facilities/managing-specific-infectious-diseases-a-to-z
6.	Specific settings and populations: additional health protection considerations	https://www.gov.uk/government/publications/health-protection-in-schools-and-other-childcare-facilities/specific-educational-settings-and-populations-additional-health-protection-considerations
7.	Children and young people settings: tools and resources (including exclusion table, posters, checklists)	https://www.gov.uk/government/publications/health- protection-in-schools-and-other-childcare- facilities/children-and-young-people-settings-tools- and-resources



Gastroenteritis / Norovirus Factsheet

Gastroenteritis

Gastroenteritis refers to a condition in which a stomach bug, usually a virus, infects the gut and gives rise to symptoms such as diarrhoea and vomiting.

Norovirus

- Norovirus, also known as the 'winter vomiting bug' is the most common cause of gastroenteritis
 in England and Wales.
- Norovirus affects people of all ages and spreads easily from person to person.
- The virus is transmitted by direct contact with an infected person, consuming contaminated food
 or water or by coming into contact with contaminated surfaces or objects.

Symptoms

Sudden onset nausea, vomiting and diarrhoea, usually 1-2 days after becoming infected. Other symptoms may include fever, a headache and body aches.

Stay at Home

Stay off the educational setting or work until you have not been sick or had diarrhoea for at least 48 hours.

Recovery

- There is no specific treatment for norovirus and most people will make a full recovery in 2-3 days without needing medical assessment or medication.
- Some people (usually the very young or elderly) may become very dehydrated and require
 hospital admission. It is therefore important to drink plenty of fluids during the illness to avoid
 dehydration.
- Those with diarrhoea and vomiting should try and avoid attending A&E or their GP surgery
 unless advised to do so by a healthcare professional as this can spread the virus to others.

Prevention

- Good hand hygiene is essential to preventing Norovirus from spreading.
- Hands should be washed thoroughly using soap and water (not alcohol gel) after using the toilet, before preparing and eating food and after contact with a sick person. Alcohol gel is not effective against norovirus, but it can be used in addition to soap and water.
- Children should be encouraged to wash their hands after using the toilet, after play, before
 eating and on arrival and departure from educational setting (to ensure they do not bring the
 infection to educational setting or take it home).
- For younger children handwashing should take place under supervision to ensure proper technique.
- Any contaminated clothing or bedding should be washed with detergent at 60°C.
- Surfaces and touch points (taps, toilet flush handle, door handles) should be cleaned and disinfected with a bleach-based cleaner.
- Stay off educational setting or work until you have not been sick or had diarrhoea for at least 48 hours.

Further Information:

Diarrhoea and vomiting - NHS (www.nhs.uk)



Best Practice: How to hand wash step by step images


Steps 3-8 should take at least 15 seconds.




Wet hands with water



Apply enough soap to cover all hand surfaces.




Rub hands palm to palm.



Right palm over the back of the other hand with interlaced fingers and vice versa.



Palm to palm with fingers interlaced.



Backs of fingers to opposing palms with fingers interlocked.



Rotational rubbing of left thumb clasped in right palm and vice versa.



Rotational rubbing, backwards and forwards with clasped fingers of right hand in left palm and vice versa.




Rinse hands with water.



Dry thoroughly with towel




Use elbow to turn off tap.



... and your hands are safe*.

Adapted from the World Health Organization/Health Protection Scotland © Crown copyright 2023. UKHSA gateway number: 2023261. *Any skin complaints should be referred to local occupational health or GP.















Hand Hygiene (e-bug.eu)