Average sturgeon weight increases to 38 pounds

Sturgeon stats

Fish

taken

1,505

661

851 464

221

520

340 262

685

718

300 1,424 21

159

117

407

39 302 783

491 842

2,238

1,251

1,001

length

42 50

Year

1955

1956

1957

1958

1959

1960

1961

1962

1963

1964

1965

1966

1967

1968

1969

1970

1971

1972

1973

1974

1975

1976

1977

1978

1979

1980

1981

1982

1983

1984

1985

1986

1988

By PAT DURKIN of the Northwestern

The average sturgeon swimming around Lake Winnebago this winter is population.

packing a few more pounds than it was one year ago.

According to the Department of Natural Resources in Oshkosh, the average weight of the 468 sturgeon speared during the February season was 38 pounds. That is three pounds more than the average in 1987.

Also, for the fourth straight year, and the sixth time in the 1980s, the average length was 55 inches.

This increase comes one year after the average weight dipped to the all-time low of 35. The DNR has kept track of the average weight since 1955. Since then, the average overall weight is about 43 pounds.

Dan Folz, area fish manager, predicted last fall that sturgeon weights would prob-

ably be higher this year. He based that sturgeon the chance to go back to their on the fact that high populations of troutperch, emerald shiners, gizzard shad, and white bass and sheepshead fry were present in the lake.

These forage fish provide plenty of

food for predatory fish such as walleyes. perch and saugers. As a result, there is less competition for the lake's redworm

> The redworms, which are lakefly larvae, make up 99 percent of the lake sturgeon's diet. When redworms are eaten heavily by other fish, there is less food available for the sturgeon.

"All last fall we kept seeing the average weight climbing for all fish," Folz said. "We assumed it was because there were so many forage fish available. If other species are getting bigger, it usually follows that the lake sturgeon will, too,"

Mark Endris, assistant area fish manager. said spearers in various parts of the lakes reported that the redworm beds appeared healthier and larger than last winter.

"With all the forage fish available, the game fish seem to be leaving the worm beds alone." he said. "That gives the

traditional food base."

Folz said the average gain in weight was still below the long-term average of

Please turn to Weight / Page D8